

All cases referred for consideration of routine surgery (asterisked) should be:

Fit for anaesthetic with Blood pressure, BMI, Fbc, u&e, LFTs, BG/urinalysis

Have measured respiratory function (where appropriate)

§ Questionnaires can be downloaded from EDICS web-site (www.edics.co.uk)

Acne for isotretinoin therapy

Routine treatments all tried

Fbc, LFTs, fasting lipids

Females not pregnant

Hip replacement*

§ Oxford hip score (>34 significant)

Knee replacement*

§ Oxford knee score (>34 significant)

Benign prostatic hypertrophy*

PSA (where relevant), u&e

§ IPSS score

Prostatic/bladder ultrasound

Flow rate (where available)

Infertility investigations

Day 21 progesterone

Day 2-4 FSH/LH

Prolactin (if amenorrhoea)

Sperm count

Cx smear

Endocervical swab for chlamydia & gonorrhoea

Rubella status

Syphilis status

Dermatology

SPCT low priority exclusions noted

Duration, size, colour, site

Unusual characteristics

Working diagnosis

Dyspepsia

BMI

Fbc, LFTs, other bloods

Results of endoscopy

Helicobacter stool test

Colonoscopy

Duration of symptoms

BMI, weight loss

Fbc, LFTs, u&e

Availability of carer

Transport requirements

Gastroscopy

BMI

Availability of carer

Transport requirements

Abnormal LFTs

BMI

Fbc, blood glucose, u&e

Serial LFTs where available (inc GGT)

Hep A/B/C status

Immunoglobulins

Autoantibodies

Alpha-1 antitrypsin

Ceruloplasmin

Ferritin

Ultrasound scan of liver

Recurrent miscarriage

Thrombophilia screen (lupus, APCR-F5)

Thyroid antibodies, ANF, anticardiolipin

Hormonal profile Day 2-7 (LH, FSH,

prolactin, SHBG, free androgen index, testosterone)

Nosebleeds

Trial of topical antibiotic cream

Suspected parotid/submandibular stone

Sialogram

Chronic rhinitis

Consider trial of intranasal steroid (spray/drops)